arXiv:1402.2361v1 [math.CO] 11 Feb 2014

An Explicit Formula for Bell Numbers in Terms of Stirling Numbers and Hypergeometric Functions

Feng Qi

Department of Mathematics, College of Science
Tianjin Polytechnic University
Tianjin City, 300387,
People's Republic of China
qifeng618@gmail.com
qifeng618@hotmail.com
qifeng618@qq.com
http://qifeng618.wordpress.com

Abstract

In the paper, the author finds an explicit formula for computing Bell numbers in terms of Kummer confluent hypergeometric functions and Stirling numbers of the second kind.

1 Introduction

In combinatorics, Bell numbers, usually denoted by B_n for $n \in \{0\} \cup \mathbb{N}$, count the number of ways a set with n elements can be partitioned into disjoint and non-empty subsets. These numbers have been studied by mathematicians since the 19th century, and their roots go back to medieval Japan, but they are named after Eric Temple Bell, who wrote about them in the 1930s. Every Bell number B_n may be generated by

$$e^{e^x-1} = \sum_{k=0}^{\infty} \mathrm{B}_k \frac{x^k}{k!} \tag{1}$$

or, equivalently, by

$$e^{e^{-x}-1} = \sum_{k=0}^{\infty} (-1)^k \mathrm{B}_k \frac{x^k}{k!}. \tag{2}$$

In combinatorics, Stirling numbers arise in a variety of combinatorics problems. They are introduced in the eighteen century by James Stirling. There are two kinds of Stirling

numbers: Stirling numbers of the first kind and Stirling numbers of the second kind. Every Stirling number of the second kind, usually denoted by $S(n,k)$, is the number of ways of partitioning a set of n elements into k nonempty subsets, may be computed by

$$S(n,k)=\frac{1}{k!}\sum_{i=0}^{k}(-1)^i\binom{k}{i}(k-i)^n, \tag{3}$$

and may be generated by

$$\frac{(e^x-1)^k}{k!}=\sum_{n=k}^{\infty}S(n,k)\frac{x^n}{n!},\quad k\in\{0\}\cup\mathbb{N}. \tag{4}$$

In the theory of special functions, the hypergeometric functions are denoted and defined by

$${}_pF_q(a_1,\dots,a_p;b_1,\dots,b_q;x)=\sum_{n=0}^{\infty}\frac{(a_1)_n\cdots(a_p)_n}{(b_1)_n\cdots(b_q)_n}\frac{x^n}{n!} \tag{5}$$

for $b_i\notin\{0,-1,-2,\dots\}$ and $p,q\in\mathbb{N}$, where $(a)_0=1$ and $(a)_n=a(a+1)\cdots(a+n-1)$ for $n\in\mathbb{N}$ and any complex number a is called the rising factorial. Specially, the series

$${}_1F_1(a;b;z)=\sum_{k=0}^{\infty}\frac{(a)_k}{(b)_k}\frac{z^k}{k!} \tag{6}$$

is called Kummer confluent hypergeometric function.

In combinatorics or number theory, it is common knowledge that Bell numbers B_n may be computed in terms of Stirling numbers of the second kind $S(n,k)$ by

$$\mathrm{B}_n=\sum_{k=1}^{n}S(n,k). \tag{7}$$

In this paper, we will find a new explicit formula for computing Bell numbers B_n in terms of Kummer confluent hypergeometric functions ${}_1F_1(k+1;2;1)$ and Stirling numbers of the second kind $S(n,k)$ as follows.

Theorem 1. *For $n\in\mathbb{N}$, Bell numbers B_n may be expressed as*

$$\mathrm{B}_n=\frac{1}{e}\sum_{k=1}^{n}(-1)^{n-k}S(n,k)k!{}_1F_1(k+1;2;1). \tag{8}$$

2 Proof of Theorem 1

We now start out to verify Theorem 1 as follows.

Among other things, Qi and Wang [5, Theorem 1.2] obtained that the function

$$H_k(z) = e^{1/z} - \sum_{m=0}^{k} \frac{1}{m!}\frac{1}{z^m} \tag{9}$$

for $k \in \{0\} \cup \mathbb{N}$ and $z \neq 0$ has the integral representation

$$H_k(z) = \frac{1}{k!(k+1)!}\int_0^\infty {}_1F_2(1; k+1, k+2; t)t^k e^{-zt}\,\mathrm{d}\,t, \quad \Re(z) > 0. \tag{10}$$

See also [3, Section 1.2] and [4, Lemma 2.1]. When $k = 0$, the integral representation (10) becomes

$$e^{1/z} = 1 + \int_0^\infty \frac{I_1\bigl(2\sqrt{t}\,\bigr)}{\sqrt{t}} e^{-zt}\,\mathrm{d}\,t, \quad \Re(z) > 0, \tag{11}$$

where $I_\nu(z)$ stands for the modified Bessel function of the first kind

$$I_\nu(z) = \sum_{k=0}^{\infty} \frac{1}{k!\Gamma(\nu + k + 1)}\left(\frac{z}{2}\right)^{2k+\nu} \tag{12}$$

for $\nu \in \mathbb{R}$ and $z \in \mathbb{C}$, see [1, p. 375, 9.6.10], and Γ represents the classical Euler gamma function which may be defined by

$$\Gamma(z) = \int_0^\infty t^{z-1}e^{-t}\,\mathrm{d}\,t, \quad \Re z > 0, \tag{13}$$

see [1, p. 255]. Replacing z by e^x in (11) gives

$$e^{1/e^x} = e^{e^{-x}} = 1 + \int_0^\infty \frac{I_1\bigl(2\sqrt{t}\,\bigr)}{\sqrt{t}} e^{-e^x t}\,\mathrm{d}\,t. \tag{14}$$

Differentiating $n \geq 1$ times with respect to x on both sides of (14) and (2) gives

$$\frac{\mathrm{d}^n\, e^{e^{-x}}}{\mathrm{d}\,x^n} = \int_0^\infty \frac{I_1\bigl(2\sqrt{t}\,\bigr)}{\sqrt{t}} \frac{\mathrm{d}^n\, e^{-e^x t}}{\mathrm{d}\,x^n}\,\mathrm{d}\,t \tag{15}$$

and

$$\frac{\mathrm{d}^n\, e^{e^{-x}}}{\mathrm{d}\,x^n} = e\sum_{k=n}^{\infty}(-1)^k \mathrm{B}_k \frac{x^{k-n}}{(k-n)!}. \tag{16}$$

From (15) and (16), it follows that

$$e\sum_{k=n}^{\infty}(-1)^k \mathrm{B}_k \frac{x^{k-n}}{(k-n)!} = \int_0^\infty \frac{I_1\bigl(2\sqrt{t}\,\bigr)}{\sqrt{t}} \frac{\mathrm{d}^n\, e^{-e^x t}}{\mathrm{d}\,x^n}\,\mathrm{d}\,t.$$

Taking $x \to 0$ in the above equation yields

$$(-1)^n e\mathrm{B}_n = \int_0^\infty \frac{I_1\bigl(2\sqrt{t}\,\bigr)}{\sqrt{t}} \lim_{x\to0} \frac{\mathrm{d}^n\, e^{-e^x t}}{\mathrm{d}\, x^n}\,\mathrm{d}\, t. \tag{17}$$

In combinatorics, Bell polynomials of the second kind, or say, the partial Bell polynomials, $\mathrm{B}_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ are defined by

$$\mathrm{B}_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1\le i\le n, \ell_i\in\mathbb{N}\\ \sum_{i=1}^n i\ell_i=n\\ \sum_{i=1}^n \ell_i=k}} \frac{n!}{\prod_{i=1}^{n-k+1}\ell_i!} \prod_{i=1}^{n-k+1}\Bigl(\frac{x_i}{i!}\Bigr)^{\ell_i} \tag{18}$$

for $n \ge k \ge 1$, see [2, p. 134, Theorem A], and satisfy

$$\mathrm{B}_{n,k}\bigl(abx_1, ab^2x_2, \dots, ab^{n-k+1}x_{n-k+1}\bigr) = a^k b^n \mathrm{B}_{n,k}(x_1, x_n, \dots, x_{n-k+1}) \tag{19}$$

and

$$\mathrm{B}_{n,k}(\overbrace{1, 1, \dots, 1}^{n-k+1}) = S(n, k), \tag{20}$$

see [2, p. 135], where a and b are any complex numbers. The well-known Faà di Bruno formula may be described in terms of Bell polynomials of the second kind $\mathrm{B}_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ by

$$\frac{\mathrm{d}^n}{\mathrm{d}\, x^n} f\circ g(x) = \sum_{k=1}^n f^{(k)}(g(x))\mathrm{B}_{n,k}\bigl(g'(x), g''(x), \dots, g^{(n-k+1)}(x)\bigr), \tag{21}$$

see [2, p. 139, Theorem C]. By Faà di Bruno formula (21) and the identities (19) and (20), we have

$$\begin{aligned}
\frac{\mathrm{d}^n\, e^{-e^x t}}{\mathrm{d}\, x^n} &= \sum_{k=1}^n e^{-e^x t}\mathrm{B}_{n,k}(\overbrace{-e^x t, -e^x t, \dots, -e^x t}^{n-k+1})\\
&= e^{-e^x t}\sum_{k=1}^n (-e^x t)^k \mathrm{B}_{n,k}(\overbrace{1, 1, \dots, 1}^{n-k+1})\\
&= e^{-e^x t}\sum_{k=1}^n (-e^x t)^k S(n, k)\\
&\to e^{-t}\sum_{k=1}^n (-t)^k S(n, k)
\end{aligned} \tag{22}$$

as $x \to 0$. Substituting (22) into (17) leads to

$$\begin{aligned}
\mathrm{B}_n &= \frac{1}{e}\sum_{k=1}^n (-1)^{n-k} S(n, k)\int_0^\infty I_1\bigl(2\sqrt{t}\,\bigr) t^{k-1/2} e^{-t}\,\mathrm{d}\, t\\
&= \frac{1}{e}\sum_{k=1}^n (-1)^{n-k} S(n, k) k!\, {}_1F_1(k+1; 2; 1).
\end{aligned}$$

The required proof is complete.

References

[1] M. Abramowitz and I. A. Stegun (Eds), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Applied Mathematics Series **55**, 10th printing, Dover Publications, 1972. 3

[2] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., 1974. 4

[3] F. Qi, Properties of modified Bessel functions and completely monotonic degrees of differences between exponential and trigamma functions, preprint, `http://arxiv.org/abs/1302.6731`. 3

[4] F. Qi and C. Berg, Complete monotonicity of a difference between the exponential and trigamma functions and properties related to a modified Bessel function, *Mediterr. J. Math.* **10** (2013), 1685–1696; Available online at `http://dx.doi.org/10.1007/s00009-013-0272-2`. 3

[5] F. Qi and S.-H. Wang, Complete monotonicity, completely monotonic degree, integral representations, and an inequality related to the exponential, trigamma, and modified Bessel functions, preprint, `http://arxiv.org/abs/1210.2012`. 3

2010 *Mathematics Subject Classification*: Primary 11B73; Secondary 33B10, 33C15.
Keywords: explicit formula, Bell number, confluent hypergeometric function of the first kind, Stirling number of the second kind.

(Concerned with sequences A000110 and A008277.)